

# Decree® 50 WDG

Fungicide



SPECIMEN

GROUP 17 FUNGICIDE

50% Water Dispersible Granule for Control of *Botrytis* in Various Greenhouse Crops, *Botrytis* Diseases in Ornamentals, Forestry Conifers and Non-Bearing Fruit Trees and Vines and for the Suppression of Powdery Mildew in Ornamentals.

**FOR AGRICULTURAL AND ORNAMENTAL HORTICULTURAL USE ONLY**  
Not for Residential Use

**Active Ingredient**

Fenhexamid† .....	50.0%
<b>Other Ingredients</b> .....	50.0%
<b>TOTAL</b> .....	100.0%

†N-(2,3-dichloro-4-hydroxyphenyl)-1-methyl-cyclohexane carboxamide

**Keep Out of Reach of Children**

## CAUTION

**NOTICE:** Read the entire label before using. Use only according to label directions.

**Before using this product, read Conditions of Sale at the end of label booklet.**

EPA Reg. No. 66330-35-67690

FPL20120425

Manufactured for: **SePRO Corporation**

11550 North Meridian Street, Suite 600 Carmel, IN 46032, U.S.A.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE:</b> Call PROSAR at 1-866-303-6952	
<b>FOR 24-HOUR CHEMICAL EMERGENCY:</b> (Spill, Leak, Fire, Exposure or Accident) Call INFOTRAC at 1-800-535-5053	

Refer to inside of label booklet for additional precautionary information and directions for use, including storage and disposal.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Socks and shoes; and
- Chemical-resistant gloves made of any waterproof material.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during the application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated; such as plants, soil, or water is: coveralls, socks, shoes and chemical resistant gloves made of waterproof material.

Carefully read and understand the Directions for Use and restrictions before applying this product.

Decree 50 WDG Fungicide is a hydroxylanilide protectant fungicide used for the control of Gray mold (*Botrytis cinerea*) on various greenhouse vegetables and for control of Gray mold or *Botrytis* blights (*Botrytis cinerea*, *Botrytis elliptica* and *Botrytis tulpae*) on ornamental crops, forestry conifers and non-bearing fruit trees and vines. When Decree 50 WDG is used to control *Botrytis* diseases, suppression of powdery mildew (*Uncinula necator*) on ornamentals crops may be achieved. Decree 50 WDG should be used in a program with other fungicides that are labeled for the control of powdery mildew. Decree 50 WDG should be used in a program with other products to provide season-long *Botrytis* protection.

Apply as directed in the *APPLICATION* section using sufficient water to obtain thorough coverage of plants. Under severe disease conditions, use maximum rates and shorter application intervals as specified on the label. Dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly.

**RESISTANCE MANAGEMENT**

Decree 50 WDG is a group 17 fungicide. While there is no indication of cross-resistance between the active ingredient in this product and other fungicides, no more than two consecutive applications should be made with Decree 50 WDG or any product containing its active ingredient, fenhexamid. After the second application, use an alternative fungicide effective in controlling the labeled diseases for two applications before reapplying the active ingredient in this product. Consult your local crop advisor for appropriate rotation recommendations. Resistance management strategies such as a blocking program, alternation or tank mixture with other fungicides used for *Botrytis* control are recommended. To avoid the development of resistant strains, do not apply Decree 50 WDG at rates lower than recommended on this label.

**COMPATIBILITY OF MIXTURES**

This product is believed to be compatible with most commonly used fungicides, insecticides, micronutrients, growth regulators and spray adjuvants. Consult specific product labels for additional information. It is always advisable to conduct a tank spray compatibility test before mixing this product with other products. To determine the physical compatibility of this product with other products, use a jar test. (Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry formulations first, then flowables, then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible.) Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. It is suggested that combinations be used on a small number of plants before treating large areas.

**PREPARATION OF SPRAY SOLUTION**

**Read *COMPATIBILITY OF MIXTURES* section prior to any tank mixing with this product.** Begin with clean equipment. Add sufficient clean water to the spray tank for one-half of the mix load. Start tank agitation. Add and properly suspend the necessary amount of product before adding any other products. Agitate to ensure thorough mixing while adding the remaining required water. A high quality wetting agent or other spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions. Decree 50 WDG should be properly suspended and diluted prior to the addition of any adjuvant. When an adjuvant is to be used with this product, SePRO recommends the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant. Consult the adjuvant label or manufacturer for crop tolerance and safety information when used with this product. Maintain agitation during mixing. Do not allow the spray mixture to stand without agitation. Apply with properly calibrated spray equipment.

**APPLICATION**

Prepare only the amount of spray solution needed to treat the area to be sprayed. Maintain agitation during application. Applications should be made immediately after the spray solution is prepared. If there is any delay in application, continue mix agitation. Do not allow the spray mixture to stand without agitation. If agitation is stopped for any reason, the spray solution must be thoroughly re-mixed prior to any further use. Rainfall or overhead irrigation within 2 hours of application may necessitate retreatment.

Proper calibration and operation of spray equipment is essential for successful spraying. Consult the spray equipment manufacturer's operator's manual, spray nozzle catalogs, and/or your crop advisor for more information.

Do not apply this product through any type of irrigation system.

For agricultural applications, apply using ground equipment only (such as boom type, air blast, hand gun or mist blower sprayer). For food crops, thorough coverage of all flowers and fruiting parts is essential for control of blossom and fruit infections caused by *Botrytis*.

For ornamental crops, thorough coverage is essential for disease control. Apply in sufficient water to obtain complete coverage of flowers, foliage and stems. Applications may be made with high volume, low volume or ultra low volume (non-thermal foggers, misters, etc.) ground equipment only. Follow the spray equipment manufacturer's directions to determine the amount of spray solution required to obtain thorough coverage.

Consult the spray equipment manufacturer's operator manual, spray nozzle catalogs and/or your crop advisor for more information. Do not apply this product using thermal application equipment.

Do not plant food crops in the treated field other than those for which the product is labeled within 30 days following the last application.

<b>CILANTRO</b>			
<b>Transplant and Greenhouse Production (except California)</b>			
<b>Transplant Production</b>			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)	<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat application after 7 days if conditions continue to favor disease development.</li> </ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)	
<b>Greenhouse Production</b>			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)	<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat application after 7 days if conditions continue to favor disease development.</li> <li>Do not apply within 3 days of harvest.</li> </ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)	
<b>Restrictions</b>			
<ul style="list-style-type: none"> <li>Do not apply in field.</li> <li>Do not make more than 2 consecutive applications. Refer to the <i>Resistance Management</i> section for rotation guidelines.</li> <li>Under conditions conducive to low or moderate disease pressure only, Decree 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb a.i.) per acre with the recommended rate of another fungicide which is registered for use on cilantro to control <i>Botrytis cinerea</i>.</li> </ul>			
<b>Transplant Production</b>			
<ul style="list-style-type: none"> <li>Apply in a full coverage protective spray.</li> <li>Do not apply more than 3.0 lbs Decree 50 WDG (1.5 lbs a.i.) per acre to transplants.</li> </ul>			
<b>Greenhouse Production</b>			
<ul style="list-style-type: none"> <li>PHI = 3 days.</li> <li>Do not apply on transplanted cilantro prior to bloom.</li> <li>Do not apply more than 3.0 lbs of Decree 50 WDG (1.5 lbs a.i.) per acre per season.</li> </ul>			

<b>CUCUMBER</b>			
<b>Transplant and Greenhouse Production (except California)</b>			
<b>Transplant Production</b>			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)	<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat at 7-10 day intervals if conditions continue to favor disease.</li> </ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)	
<b>Greenhouse Production</b>			
Disease	Application Rate		Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)	<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat at 7-10 day intervals if conditions continue to favor disease.</li> <li>May be applied up to and including the day of harvest.</li> </ul>
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)	

(continued)

<b>CUCUMBER (Cont.)</b> <b>Transplant and Greenhouse Production (except California)</b>		
<b>Restrictions</b>		
<ul style="list-style-type: none"> <li>Do not apply in field.</li> <li>Do not make more than 2 consecutive applications. Refer to the <i>Resistance Management</i> section for rotation guidelines.</li> <li>Under conditions conducive to low or moderate disease pressure only, Decree 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb a.i.) per acre with the recommended rate of another fungicide which is registered for use on cucumber to control <i>Botrytis cinerea</i>.</li> </ul>		
<b>Transplant Production</b>		
<ul style="list-style-type: none"> <li>Apply in a full coverage protective spray.</li> <li>Do not apply more than 6.0 lbs Decree 50 WDG (3.0 lbs a.i.) per acre to transplants.</li> </ul>		
<b>Greenhouse Production</b>		
<ul style="list-style-type: none"> <li>PHI = 0 day.</li> <li>Do not apply more than 6.0 lbs of Decree 50 WDG (3.0 lbs a.i.) per acre per season.</li> </ul>		

<b>FRUITING VEGETABLES (except Tomatoes)</b> <b>Eggplant; Ground Cherry; Pepino; Pepper (Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper); and Tomatillo</b> <b>Transplant and Greenhouse Production (except California)</b>		
<b>Transplant Production</b>		
Disease	Application Rate	Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)
<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat at 7-10 day intervals if conditions continue to favor disease.</li> </ul>		
<b>Greenhouse Production</b>		
Disease	Application Rate	Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)
<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat at 7-10 day intervals if conditions continue to favor disease.</li> <li>May be applied up to and including the day of harvest.</li> </ul>		
<b>Restrictions</b>		
<ul style="list-style-type: none"> <li>Do not apply in field.</li> <li>Do not make more than 2 consecutive applications. Refer to the <i>Resistance Management</i> section for rotation guidelines.</li> <li>Under conditions conducive to low or moderate disease pressure only, Decree 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb a.i.) per acre with the recommended rate of another fungicide which is registered for use on fruiting vegetables to control <i>Botrytis cinerea</i>.</li> </ul>		
<b>Transplant Production</b>		
<ul style="list-style-type: none"> <li>Apply in a full coverage protective spray.</li> <li>Do not apply more than 6.0 lbs Decree 50 WDG (3.0 lbs a.i.) per acre to transplants.</li> </ul>		
<b>Greenhouse Production</b>		
<ul style="list-style-type: none"> <li>PHI = 0 day.</li> <li>Do not apply more than 6.0 lbs of Decree 50 WDG (3.0 lbs a.i.) per acre per season.</li> </ul>		

<b>TOMATOES</b> <b>Transplant and Greenhouse Production</b>		
<b>Transplant Production</b>		
Disease	Application Rate	Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)
<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat at 7-10 day intervals if conditions continue to favor disease.</li> </ul>		
<b>Greenhouse Production</b>		
Disease	Application Rate	Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)
<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat at 7-10 day intervals if conditions continue to favor disease.</li> <li>May be applied up to and including the day of harvest.</li> </ul>		
<b>Restrictions</b>		
<ul style="list-style-type: none"> <li>Do not apply in field.</li> <li>Do not make more than 2 consecutive applications. Refer to the <i>Resistance Management</i> section for rotation guidelines.</li> <li>Under conditions conducive to low or moderate disease pressure only, Decree 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb a.i.) per acre with the recommended rate of another fungicide which is registered for use on tomatoes to control <i>Botrytis cinerea</i>.</li> </ul>		
<b>Transplant Production</b>		
<ul style="list-style-type: none"> <li>Apply in a full coverage protective spray.</li> <li>Do not apply more than 6.0 lbs Decree 50 WDG (3.0 lbs a.i.) per acre to transplants.</li> </ul>		
<b>Greenhouse Production</b>		
<ul style="list-style-type: none"> <li>PHI = 0 day.</li> <li>Do not apply more than 6.0 lbs of Decree 50 WDG (3.0 lbs a.i.) per acre per season.</li> </ul>		

<b>LEAFY GREENS (except Spinach)</b> <b>Amaranth; Arugula; Chervil; Chrysanthemum, Edible-leaved and Garland; Corn Salad; Cress, Garden and Upland; Dandelion; Dock; Endive; Lettuce; Orach; Parsley; Purslane, Garden and Winter; Radicchio (Red chicory)</b> <b>Transplant and Greenhouse Production (except California)</b>		
<b>Transplant Production</b>		
Disease	Application Rate	Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)
<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat application after 7 days if conditions continue to favor disease.</li> </ul>		
<b>Greenhouse Production</b>		
Disease	Application Rate	Application Timing
Gray mold ( <i>Botrytis cinerea</i> )	Stand-alone	1.5 lbs/A (0.75 lb a.i./A)
	Tank-mix	1.0 - 1.5 lbs/A (0.5 - 0.75 lb a.i./A)
<ul style="list-style-type: none"> <li>Begin applications when conditions favor disease development.</li> <li>Repeat application after 7 days if conditions continue to favor disease.</li> <li>Do not apply within 3 days of harvest.</li> </ul>		

(continued)

### LEAFY GREENS (except Spinach) Cont.

Amaranth; Arugula; Chervil; Chrysanthemum, Edible-leaved and Garland; Corn Salad; Cress, Garden and Upland; Dandelion; Dock; Endive; Lettuce; Orach; Parsley; Purslane, Garden and Winter; Radicchio (Red chicory) Transplant and Greenhouse Production (except California)

#### Restrictions

- Do not apply in field.
- Do not make more than 2 consecutive applications. Refer to the *Resistance Management* section for rotation guidelines.
- Under conditions conducive to low or moderate disease pressure only, Decree 50 WDG may be applied in a tank-mix at a rate of 1.0 lb (0.5 lb a.i.) per acre with the recommended rate of another fungicide which is registered for use on leafy greens to control *Botrytis cinerea*.

#### Transplant Production

- Apply in a full coverage protective spray.
- Do not apply more than 3.0 lbs Decree 50 WDG (1.5 lbs a.i.) per acre to transplants.

#### Greenhouse Production

- PHI = 3 days.
- Do not apply more than 3.0 lbs of Decree 50 WDG (1.5 lbs a.i.) per acre per crop.

### OUTDOOR-, GREENHOUSE- AND NURSERY-GROWN ORNAMENTALS, FORESTRY CONIFERS AND NON-BEARING FRUIT TREES AND VINES

This product, when applied as a foliar spray, has been shown to effectively protect plants from *Botrytis cinerea*, *Botrytis elliptica* and *Botrytis tulipae* and stop sporulation of the fungus if an infection has occurred. Also, when this product is used to control *Botrytis*, suppression of powdery mildew on the treated crop may be achieved. These directions are based upon the results of product testing programs.

This product has been shown to be safe and effective in a limited number of research trials on certain varieties or cultivars of the plant types listed. Users should conduct small scale tests under local growing conditions prior to wide-scale use.

African Violet	Iris	Rose
Crabapple, Flowering <sup>1</sup>	Lily (Asiatic/Oriental)	Spathiphyllum
Daffodil	Lisianthus	Snagdragon
Douglas Fir	Pansy	Tulip
Fuchsia	Petunia	Western Hemlock
Geranium	Plum, Flowering <sup>1</sup>	
Gerbera Daisy	Poinsettia <sup>2</sup>	
Impatiens	Ranunculus	

<sup>1</sup> Not for food or feed purposes.

<sup>2</sup> Due to the sensitivity of mature Poinsettia bracts and varying local environmental conditions, users should conduct preliminary tests as suggested above.

#### High Volume Sprays

Application Rate	Disease Pressure	Spray Interval
0.75 lb product per 100 gal.	Conditions favor disease development; no disease present.	7 to 14 Days
1.0 to 1.5 lbs product per 100 gal.	Conditions favor disease development; disease present.	7 Days

#### Low Volume / Ultra Low Volume Sprays

Application Rate	Disease Pressure	Spray Interval
0.75 lb product per 100 gal.	Conditions favor disease development; no disease present.	7 to 14 Days
1.0 to 1.5 lbs product per 100 gal.	Conditions favor disease development; disease present.	7 Days

For small application equipment, the scoop attached to the container is provided for ease in measuring the amount of this product to use per 1 gallon of water. Apply 1 to 2 level scoops per 1 gallon of water. One scoop equals 0.12 ounces or 3.4 grams or approximately one rounded teaspoon of this product.

#### POST-HARVEST APPLICATIONS

For use on cut flowers, rose budwood, and bare-root nursery stock when applied as a spray. Apply Decree 50 WDG to prevent *Botrytis* diseases (Gray mold) in cold storage or transit. Apply after harvest and grading prior to cold storage. Apply 1.0 - 1.5 pounds of Decree 50 WDG per 100 gallons of spray.

#### LIMITATIONS

Do not make more than two consecutive applications of this product. After the second application, use an alternative fungicide effective in controlling *Botrytis spp.* for two applications before reapplying the active ingredient in this product. Consult your local Horticultural Advisor for the most appropriate alternative products.

#### Ornamentals and Other Nursery Crops Grown Outdoors

Do not apply more than 6.0 pounds of product per acre per season (3.0 lbs a.i./A/season).

#### Ornamentals and Other Nursery Crops Grown in Greenhouses

Do not apply more than 0.14 pounds product per 1,000 square feet per crop cycle (equal to 2.2 oz. product per 1,000 square feet per crop cycle).

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a secure dry place away from excessive heat. Do not store near food or feed. Store in original container only. To close package, replace and tighten cap to form an airtight seal.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

#### Container Disposal (Plastic):

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or (if allowed by State and local authorities) by burning. If burned stay out of smoke. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

#### CONDITIONS OF SALE

1. SePRO Corporation (SePRO) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the Directions for Use under normal conditions of use.
2. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to SePRO. **SEPRO DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY. TO THE EXTENT PERMITTED BY LAW, SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM OR IN CONNECTION WITH THE MANUFACTURE, SALE, DELIVERY, USE, HANDLING OR STORAGE OF THIS PRODUCT. TO THE EXTENT PERMITTED BY LAW, SELLER'S LIABILITY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. SePRO DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**
3. Critical and unforeseeable factors beyond SePRO's control prevent SePRO from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which the product is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated on the label. Except as stated in 1 above, to the extent permitted by law, by purchasing, accepting and using this product, the buyer and user acknowledge and assume all risks and liabilities resulting from handling, storage, and use of this product.
4. For species and their varieties or cultivars not specifically listed on the label, the user assumes all risk from phytotoxicity or unacceptable residue or disease control.

© Copyright 2013 SePRO Corporation

Fenhexamid was invented and is produced by Bayer AG, Germany. Decree is a registered trademark of Arysta LifeScience North America Corporation